

## Resistance against insect pests by phenolics and their derivative compounds

S. A. DAR, <sup>1</sup>A. B. WANI, MUSHTAQ, A. GANIE, A.A. KANDOO AND <sup>2</sup>M.Y. WANI

Department of Entomology; <sup>1</sup> Department of Biotechnology; <sup>2</sup> Department of Sericulture  
Sher-e-Kashmir University of Agricultural Science and Technology,  
Shalimar, Srinagar

Received : 19-01-2017; Revised : 22-08-2017; Accepted : 25-08-2017

### ABSTRACT

The chemical nature of plant phenolics may vary from a simple monomeric unit to highly polymerized structures of varying proportions of monomeric aglycon units. Phenolics are aromatic benzene ring compounds with one or more hydroxyl groups produced by the plants for protection against biotic stresses. The functions of phenolic compounds in plant physiology is difficult to overestimate. Since, they provide structural integrity and scaffolding support to plants. Importantly, phenolic phytoalexins secreted by wounded or otherwise perturbed plants, repel or kill many insect pests. Insects represent adaptive characters that have been subjected to natural selection during evolution. Plants synthesize a greater array of secondary compounds than animals because they cannot rely on physical mobility to escape their predators and have therefore evolved a chemical defence against such predators. This article, review the role of plant phenols and polyphenols, interactions and their concern to resistance mechanisms against biotic stresses.

**Keywords:** Biotic, interaction, phenolics, resistance.

Plants have prepared various secondary metabolites for resistance against biotic agents like the resistance against *Leucinodes orbonalis* by phenolic compounds of *Solanum melongena* (Dar *et al.*, 2014). Similarly the defensive plant proteins play a significant role in direct interactions between herbivores and plants, the role of non-protein secondary metabolites is huge. The term 'secondary' is used to contrast them with metabolites that are directly involved in growth, development or reproduction, although it is not always possible to determine the precise physiological role of each metabolite. Across the plant kingdom, there is a staggering diversity of secondary metabolites, and they can be distinct for small phylogenetic groups. Many secondary metabolites have been implicated in plant defences or are stress-related. Despite the rich diversity of secondary metabolites, their biosynthetic origins allow them to be classified into three basal groups : phenolics; isoprenoids; and N-containing compounds.

Kant *et al.* (2015) reviewed the plants to preserve their fitness when attacked by herbivores, it can employ complex strategies that include reallocation of resources and the production of defensive metabolites and structures, which can be either prefabricated or be produced only upon attack.

### Phenolics

Phenolic compounds are secondary metabolites which are synthesized in plants and possess biological properties like antefedent (Dar, 2012), antioxidant, antiapoptosis, anti-aging, anticarcinogen, anti-inflammation, anti-atherosclerosis, cardiovascular

protection, inhibition of angiogenesis and cell proliferation activity. Phenolics consist of an aromatic ring with one or more hydroxyl groups. Two pathways are responsible for the majority of plant phenolics: (1) the phenylpropanoid pathway (Cheyner *et al.*, 2013), which converts the aromatic amino acid phenylalanine into phenolics. Phenylpropanoid are a diverse chemical class with immense health benefits that are biosynthesized from the aromatic amino acid L-phenylalanine (Dwivedi *et al.*, 2016), and (2) the acetate/malonate (polyketide) pathway (Quideau *et al.*, 2011). Single phenolics can be polymerized to form polyphenols and both can be subjected to additional modifications, giving rise to a vast quantity (>9000) of chemically diverse metabolites, which include benzoquinones, phenolic acids, coumarins, flavonoids, lignins and tannins (Balasundram *et al.*, 2006). Most of these biological actions have been attributed to their intrinsic reducing capabilities (Han, 2007). Rich dietary sources of phenolics include fruits, tea, coffee, cocoa and processed foods derived from these, such as wine. At high levels, and in particular when sugar levels are low, phenols impart an astringency, bitterness and color to foods like brinjal. Phenolics in grapes and wines include many different substances: phenolic acids (e.g., hydroxybenzoic acids such as gallic acid, hydroxycinnamic acids found in grape juice), three classes of flavonoids found in the skins and seeds (red anthocyanins, the flavonols and the abundant flavan-3-ols, comprise the monomeric catechins), oligomeric proanthocyanidins, and polymeric condensed tannins. Plant foods like brinjal (Dar, 2012) have phenolic

compounds, which affect their appearance, taste, odour and oxidative stability. In cereal grains, these compounds are located mainly in the pericarp (Naczka *et al.*, 2004). The major phenolic acids in cereals are ferulic and p-coumaric acids (Holtekjolen *et al.*, 2006 and Mattila *et al.*, 2005). Anthocyanins are water-soluble pigments mostly studied in cereals (Yao *et al.*, 2004).

Phenolics have various functions in primary metabolism; for instance, they protect plants from UV radiation (Landry *et al.*, 1995) or form, as lignins, an integral part of the secondary cell wall in vascular plants (Boerjan *et al.*, 2003). Furthermore, flavonoids are crucial for reproduction because they are required for pollen development (Van der Meer *et al.*, 1992) and provide many of the visual and volatile cues used by flowers (Hoballah *et al.*, 2007) and fruits (Jaakola *et al.*, 2002) to attract pollinators and seed dispersers, respectively. Plants use phenolics to resist attacks from herbivores because of their deterrent (Kessler *et al.*, 2012b) and toxic (Lindroth and Peterson, 1988) nature. Hence, they are often constitutively present at or near the cell surface or stored as inactive compounds away from activating enzymes e.g. in vacuoles or specialized cells, or bound to the cell wall (Pourcel *et al.*, 2007).

### **Plants synthesize phenolics**

Phenolics are often produced and accumulated in the subepidermal layers of plant tissues exposed to stress and pathogen attack (Cle *et al.*, 2008). The concentration of a particular phenolic compound within a plant tissue is dependent on season and may also vary at different stages of growth and development (Ozyigit *et al.*, 2007; Dar, 2012). Several internal and external factors, including trauma, wounding, drought and pathogen attack, affect the synthesis and accumulation of phenolics (Kefeli *et al.*, 2003). Furthermore, the biosynthesis of phenolics in chloroplasts and their accumulation in vacuoles are enhanced on exposure to light (Kefeli *et al.*, 2003). Photo-inhibition, as well as nutrient stresses, such as deficiencies in nitrogen, phosphate, potassium, sulphur, magnesium, boron and iron, also triggers the synthesis of phenyl propanoid compounds in some plant species, which include members of the flavonoid biosynthetic pathway (Balasundram *et al.*, 2006).

### **Phenolics for defence**

Phenolics serve a dual function of both repelling (Dar, 2012; Dar *et al.*, 2014) and attracting different organisms in the plant's surroundings, while as sugars do reverse of it (Fig 1). They act as protective agents, inhibitors, natural animal toxicants and pesticides against invading herbivores, nematodes, phytophagous insects and fungal and bacterial pathogens (Lattanzio *et al.*, 2006).

In brinjal the seed ring contains high content of phenolic compounds, so infestation by the *L. orbonalis* is least in that region, while as in pulp the concentration of the phenolics is less and sugars are more so attack is accordingly more. The thickness of the pulp is more and the pest remains continue to feed on it till several larval stages (4 stages). However, the pest feed less in the seed ring and move back to the pulp. Provide information in relation to other crops of same family at least to trace upon the trend.

### **Tannins**


Plants synthesized different types of phenolics with diverse functions that they perform against biotic stresses. In KSUAST-K, Srinager the chemical constituent namely lignin were observed to have exhibited a clear correlation negative with the *L. orbonalis* incidence. Genotypes, Shalimar Brinjal Hybrid-1 and Shalimar Brinjal Hybrid-2 showed minimum concentration of lignin. Varieties, Brinjal -85 were found least attacked by the *L. orbonalis* and recorded the shoot damage of 2.34 and total moisture, and lignin content of 78.72per cent and 34.24 mg/100g dry wt. respectively. In brinjal highest lignin and crude fiber content were observed in cv. Brinjal-85 and lowest in Shalimar Brinjal Hybrid-1 (Dar *et al.*, 2014). What about other crops?

### **Allelochemicals and plant competition**

In different families of plants various types of phenolics found are as: Umbelliferone, p-hydroxybenzoic acid, vanillyl alcohol and isoflavones. Chemoattractants in Rhizobium Sinapinic acid, syringic acid, ethylsyringamide, propylsyringamide, carbethoxyethylsyringamide, parahydroxybenzoate, ferulic acid *vir* gene inducers. In Agrobacterium, Vanillyl alcohol, bromo acetosyringone

Inhibitors of *vir* gene induction in Agrobacterium: Acetosyringone, a-hydroxyacetosyringone, p-Hydroxybenzoate. Chemoattractants in Agrobacterium and Rhizobium, and *vir* gene inducers in Agrobacterium Salicylic acid Quorum quencher in Agrobacterium Hydroquinones.

Allelochemicals like Coumarins, xanthones, anthocyanidins are phenolic derivatives and are most common in plants. Phenolics act as determinants of colour and attractants of pollinators in plants. Caffeic acid *vir* gene inducer in Agrobacterium 3, 4-Dihydroxybenzoic acid Protocatechuic acid, b-resorcylic acid, protocatechuate, p-resorcyate, catechol *vir* gene inducer in Agrobacterium Chlorogenic acid. Precursor for lignin and suberin synthesis in plants are Lignin, tannins and suberins. Further phenolics act like the structural components of plant cells.



**Fig.1: Relationship between the *Leucinodes orbonalis* (Guenee) infestation and phenolic and sugars content in brinjal shoot and fruits at different phenological stages of vegetative and reproductive growth (Showket, 2012).**



**Fig.2: Infestation of the *Leucinodes orbonalis* to pulp of the brinjal fruit (Showket, 2012).**




Figure above showed the response of herbivours towards the defensive secondary metabolites.

**Fig.3: The insect resistance response shown diagrammatically**

### Catechins plant defence

Flavonoids, flavonols, flavones, genistein, daidzein, O-acetyl daidzein, 6-O-malonylgenistin, 6-O-malonyl daidzin, glycitin, 6-O-malonylglycitin nod gene inducers in *Rhizobium Apigenin*, naringenin, luteolin Chemoattractants in *Agrobacterium* and *Rhizobium*, and nod gene inducers in *Rhizobium Gallate*, gallic acid, pyrogallic acid, syringic acid, kaempferol vir gene inducers in *Agrobacterium* Flavanones, quercetin nod gene inducers in *Rhizobium* Isoflavonoids. Chemoattractants and nod gene inducers in *Rhizobium Cajanin*, medicarpin, glyceoline, rotenone, coumestrol, phaseolin, phaseolinin, limonoids, tannins, flavonoids Phytoalexins, phytoanticipins and nematocides in plant defence acid.

### Center to defense mechanism

Against microbial invaders. Phenolics are synthesized when plant pattern recognition receptors recognize potential pathogens by conserved pathogen-associated molecular patterns (PAMPs), leading to PAMP-triggered immunity. As a result, the progress of the infection is restricted long before the pathogen gains complete hold of the plant.

### How herbivores cope with plant defences

Plants and herbivores have coevolved for over 400 million years. While plants have evolved signalling networks to regulate induced defences and diversity in their palette of secondary metabolites (Dar, 2016a), herbivores have been under pressure to evade defences (Alba *et al.*, 2011). Hence, behavioural adaptations have evolved that allow herbivores to avoid defended plant tissues (Dar, 2016b; Paschold *et al.*, 2007; Shroff *et al.*, 2008; Perkins *et al.*, 2013) as much as possible or to dismantle defensive structures such as trichomes (Cardoso, 2008) and latex channels (Rodrigues *et al.*, 2010). However, herbivores have also evolved a variety of mechanisms to cope with deterrent substances produced by their host plants. Given the enormous economic impact of herbivore resistance to agrochemicals, a large part of our knowledge of adaptations to xenobiotics comes from the field of pesticide resistance (Despres *et al.*, 2007). However, the mechanisms at play are similar to those that enable them to resist defensive phytotoxins, and a functional overlap between adaptation to agrochemicals and to phytotoxins has been suggested (Dermauw *et al.*, 2013).

### Phenolics—molecules for cross-talk in the rhizosphere

The plant rhizosphere is a dynamic ecosystem of different species, representing flora, fauna and microbes that interact with each other in a variety of

complex reactions. These interactions are mainly governed by a diverse array of phenolics exuded by the growing roots of plants, together with a host of other chemicals (Bais *et al.*, 2006). The functions performed by phenolics in the plant rhizosphere have been aptly termed the 'rhizosphere effect' (Hiltner, 1904). The root exudates generally include ions, free oxygen, water, enzymes, mucilage, a number of carbon-containing primary and secondary metabolites and, most importantly, plant phenolics. The released phenolics differ from species to species and also with time, space and location. Their concentrations in the soil range from 2.1 to 4.4 per cent in monocotyledonous plants and 0.1 to 0.6 per cent in dicotyledonous plants. Phenolics trigger redox reactions in soils and selectively influence the growth of soil microorganisms that colonize the rhizosphere. Microorganisms break down phenolics into elements that contribute towards the mineralization of soil nitrogen and the formation of humus (Halvorson *et al.*, 2009). The phenolics chelate metals and improve soil porosity, providing active absorption sites and increasing the mobility and bioavailability of elements, such as potassium, calcium, magnesium, copper, zinc, manganese, molybdenum, iron and boron, for plant roots (Seneviratne and Jayasingh, 2003). Although plants exude different phenolic compounds that are toxic to most microorganisms, *Agrobacterium* and *Rhizobium* have evolved mechanisms to counteract, nullify and even utilize these defences for their own advantage (Hartmann *et al.*, 2009). The most significant of these mechanisms that involve phenolics are: (i) chemotaxis; (ii) activation of the bacterial nodulation (nod) and virulence (*vir*) gene networks; (iii) xenobiotic detoxification; and (iv) quorum signalling.

### Chemotaxis

As in several other soil bacteria, phenolics play a pivotal role in the chemotactic responses of *Agrobacterium* and *Rhizobium* in their search for growth substrates and hosts. These serve as excellent models for signal transduction and plant–microbe interactions (Samac and Graham, 2007). Diverse plant phenolic compounds with varying substitution patterns determine the chemotactic movement of *Agrobacterium* or *Rhizobium* across chemical gradients towards higher levels of potential nutrients and lower levels of inhibitors. Although acetosyringone and umbelliferone are inhibitors of nod gene inducers, *R. leguminosarum* exhibits an exaggerated chemotactic response to high concentrations of phenolic compounds (Aguilar *et al.*, 1988). Probably, the complex nature of the root exudates of different plants in the rhizosphere (Djordjevic *et al.*, 1987) necessitates such negative regulation by some phenolics. For example, it might be required to prevent

competing rhizobia from targeting the same host plant and also for creating a favourable ecological niche for each of the species, therefore preventing nodule initiation in the vicinity of clover root tips by umbelliferone, is a good example of such negative rhizospheric interactions (Djordjevic *et al.*, 1987). Acetosyringone and hydroxyacetosyringone exuded from plant wounds are potent chemoattractants at very low concentrations, and they also act to induce the *vir* genes of *Agrobacterium* (Escobar and Dandekar, 2003). In addition to acetosyringone, a number of other phenolics and sugars are effective as chemoattractants (Brencic and Winans, 2005; Palmer *et al.*, 2004).

#### Activation of the bacterial nod and vir gene networks

Microbial gene expression following chemotaxis is influenced by a diverse array of substituted plant phenols. These plant-derived signals are termed ‘host recognition factors’ or ‘xenognosins’ (Campbell *et al.*, 2000). In both the symbiotic *Rhizobium* and pathogenic *Agrobacterium*, the same phenolics that act as chemoattractants may also regulate the expression of nod and vir genes, respectively.

#### Detoxification and biotransformation

Phenolics act as antimicrobial compounds because of their ability to disrupt nonspecifically the structural integrity of bacterial membranes and to inhibit specifically bacterial enzymes involved in electron transport (Hirsch *et al.*, 2003). It is natural that bacteria would counteract or even nullify these toxic compounds. Many bacteria are endowed with the ability to degrade and utilize toxic phenols as a source of carbon (Dua *et al.*, 2002; Lovely, 2003).

#### Quorum signalling for attaining infection

An important mechanism by which members of Rhizobiaceae monitor their environment is quorum sensing (Bjarnsholt and Givskov, 2007). The production, release and sensing of homoserine lactones (HSLs) or their acylated forms (AHL) are important for quorum sensing (Steidle *et al.*, 2001). Quorum sensing allows bacterial cell–cell communication and promotes an advantageous lifestyle for both the survival and maintenance of pathogenic or symbiotic relationships within a range of environmental niches (Joint *et al.*, 2002). In quorum sensing, cell density-dependent regulation of gene expression enables bacteria to coordinate certain adaptive processes that cannot be performed by an individual microbe.

#### How plants overcome insect pests

Quorum sensing helps rhizobia to synchronize themselves to phenolic signals on a population-wide scale and to function as multicellular organisms for successful symbiosis

Plants recognize herbivores by their molecular patterns or their elicitors. It is hypothesized that polyphosphoinositides generated at the plasma membrane play an important role as second messengers, just as they do during pathogenesis (Munnik and Nielsen, 2011). The most rapid measured responses are ion (*e.g.* Ca<sup>2+</sup> and K<sup>+</sup>) fluxes across the plasma membrane, followed by changes in the plasma membrane potential. Subsequently, a protein kinase cascade can activate the production of ROS such as hydrogen peroxide by activating an NADPH-dependent oxidase. Hydrogen peroxide can have a direct effect on herbivores or enter the cell, thus changing its redox status. The rapid increase in cytosolic Ca<sup>2+</sup> can also give rise to increased nitric oxide-mediated processes that precede the upregulation of JA levels (Zebelo and Maffei, 2015). These responses occur not only locally, but also in unattacked neighbouring cells and in distal tissues. Herbivory, the application of oral secretions to wounded leaves and aphid probing have been shown to give rise to membrane depolarization due to an electrochemical gradient between the interior and the exterior of the attacked plants cells. This membrane depolarization can travel with a speed of up to 40cm s<sup>-1</sup> through the entire plant and mutant plants with attenuated wound-induced surface potential changes exhibit a reduced JA response in distal leaves (Mousavi *et al.*, 2013).

#### Summery and conclusion

Finally, it is appealing to think of attack suppression as a primitive adaptation of the plants. However, some studies suggest this may not always be the case. Therefore generally it has negative and less efficient consequences also. In a plant–herbivore context such ‘mafia behaviour’ would mean that herbivores increase feeding upon experiencing induced plant defence. Although empirical studies of mafia behaviour are scarce, the costs of mounting an induced defence response under certain circumstances can be higher than its benefit. For example, when exposed to herbivores, *N. attenuata* plants typically initiate a JA-induced accumulation of nicotine, but the accumulation of nicotine is costly and decreases the competitive ability of the plant. Hence, when larvae of the nicotine-tolerant herbivore *M. sexta* feed on *N. attenuata*, the plant responds by downregulating the accumulation of nicotine. Therefore it is argued that this response likely reflects an adaptation of the plant to shut down an inefficient and costly defence response, because using this energy to compete with conspecifics may be more rewarding.

### Acknowledgement

Authors are thankful to Department of Science and Technology (DST), New Delhi, India for providing financial assistance. Thanks are also to SKUAST-K, Srinager for providing the computer and internet facilities.

### REFERENCES :

- Aguilar, J.M.M., Ashby, A.M., Richards, A.J.M., Loake, G.J., Watson, M.D. and Shaw, C.H. 1988. Chemotaxis of *Rhizobium leguminosarum* biovar phaseoli towards flavonoid inducers of the symbiotic nodulation genes. *J. Gen. Microbiol.* **134**: 2741-46
- Alba JM, Glas, J.J, Schimmel B.C.J, and Kant MR. 2011 . Avoidance and suppression of plant defenses by herbivores and pathogens. *J.Pl.Interactions* **6**: 221-27
- Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM .2006. The role of root exudates in rhizosphere interactions with plants and other organisms. *Ann. Rev Pl. Biol.* **57** : 233-66
- Balasundram, N., Sundram, K. and Samman, S. 2006. Phenolic compounds in plant and agri-industrial byproducts: antioxidant activity, occurrence, and potential uses. *Food Chem*, **99**:191-203.
- Bjarnsholt, T. and Givskov, M. 2007. Quorum-sensing blockade as a strategy for enhancing host defenses against bacterial pathogens. *Philos. Trans. R. Soc. B*, **362**:1213-22
- Brencic, A. and Winans, S.C. 2005. Detection of and response to signals involved in host-microbe interactions by plant-associated bacteria. *Microbiol. Mol. Biol. Rev.* **69**:155-94
- Bringmann, M., Li, E., Sampathkumar, A., Kocabek, T., Hauser, M.T. and Persson, S. 2012. POM-POM2/cellulose synthase interacting 1 is essential for the functional association of cellulose synthase and microtubules in Arabidopsis. *Pl. Cell*, **24**: 163-77
- Campbell, A.M., Tok, J.B., Zhang, J., Wang, Y., Stein, M., Lynn, D.G. and Binns, A.N. 2000. Xenogonin sensing in virulence: is there a phenol receptor in *Agrobacterium tumefaciens*? *Chem. Biol.* **7**, 65-76
- Cardoso MZ (2008) Herbivore handling of a Plants trichome: The case of *Heliconius charithonia* (L.) (Lepidoptera: Nymphalidae) and *Passiõora lobata* (Killip) Hutch. (Passiõoraceae). *Neotrop Ento.*, **37** : 247-52
- Cheynier, V., Comte, G., Davies, K. M., Lattanzio, V., and Martens, S. 2013 . Plant phenolics: recent advances on their biosynthesis, genetics, and ecophysiology. *Pl. Physiol. Biochem.* **72**:1-20.
- Clé, C., Hill, L.M., Niggeweg, R., Martin, C.R., Guisez, Y., Prinsen, E. and Jansen, M.A.K. 2008. Modulation of chlorogenic acid biosynthesis in *Solanum lycopersicum*; consequences for phenolic accumulation and UV-tolerance. *Phytochem.*, **69**: 2149-56.
- Dar, S.A., A. R. Wani., R.K. Nehru., S. H. Mir and M. J. Iqbal. 2014. Physio-chemical characteristics of brinjal genotypes imparting tolerance to brinjal shoot and fruit borer (*Leucinodes orbonalis*) under field conditions of Kashmir (India). *Eco. Env. Cons.*, **20** (4): 209-15.
- Dar, S.A. 2012 . Screening of brinjal genotypes/ varieties against brinjal shoot and fruit borer. *M.Sc. Thesis.* SKUAST, Srinager, Kashmir
- Dar, S.A. 2016 . Enzymatic adaptation in Insect pests to host Biochemicals. *Indian J. Expt.Biol.* (accepted)
- Dar, S.A. 2016 . Insect Plant Interaction-Signaling. *Indian J. Biochem. Biophy.* (In press).
- Dermauw W, Wybouw N, and Rombauts, S. 2013. A link between host plant adaptation and pesticide resistance in the polyphagous spider mite *Tetranychus urticae*. *Proc. Nat. Acad. Sci. USA.*
- Despres L, David J, and Gallet C. 2007. The evolutionary ecology of insect resistance to plant chemicals. *Trends Eco. Evol.* **22**: 298-307
- Djordjevic, M.A., Gabriel, D.W. and Rolfe, B.G. 1987. *Rhizobium* – the reined parasite of legumes. *Ann. Rev. Phytopathol.* **25**: 145-68
- Dua, M., Singh, A., Sethunathan, N. and Johri, A.K. 2002. Biotechnol- ogy and bioremediation: successes and limitations. *Appl. Microbiol. Biotechnol.* **59**: 143-52
- Dwivedi, S. L., Upadhyaya, H. D and Chung, M. 2016. Exploiting Phenylpropanoid Derivatives to Enhance the Nutraceutical Values of Cereals and Legumes. *Front. Pl. Sci.*, **7**: 763
- Escobar, M.A. and Dandekar, A.M. 2003. *Agrobacterium tumefaciens* as an agent of disease. *Trends Pl. Sci.* **8**, 380-86
- Halvorson, J.J., Gonzalez, J.M., Hagerman, A.E. and Smith, J.L. 2009. Sorption of tannin and related phenolic compounds and effects on soluble-N in soil. *Soil Biol. Biochem.* **41**: 2002-10
- Halvorson, J.J., Gonzalez, J.M., Hagerman, A.E. and Smith, J.L. 2009. Sorption of tannin and related phenolic compounds and effects on soluble-N in soil. *Soil Biol. Biochem.* **41**: 2002-10
- Han, G., X. Han, T. and Shen, H. Lou. 2007 . Dietary polyphenols and their biological significance. *Int. J. Mole. Sci.*, **8**:950-88.
- Hiltner L.(1904). Uber neue erfahrungen und probleme auf dem gebiete der bodenbakteriologie. *Arbeiten der Deutschen Landwirtschaft Gesellschaft*, **98**: 59-78.
- Hirsch, A.M., Bauer, W.D., Bird, D.M., Cullimore, J., Tyler, B. and Yoder, J.I. 2003. Molecular signals and receptors: controlling rhizo- sphere interactions between plants and other organisms. *Ecology*, **84**.

- Holtekjolen, A., K., Kinitz, C. and Knutsen, S., H., 2006. Flavanol and bound phenolic acid contents in different barley varieties. *J Agric. Food Chem.* **54**: 2253
- Joint, I., Trait, K., Callow, M.E., Callow, J.A., Milton, D., Williams P. and Cámara, M. 2002. Cell-to-cell communication across the prokaryote–eukaryote boundary. *Science*, **298**:1207.
- Kefeli VI, Kalevitch MV, and Borsari B .2003. Phenolic cycle in plants and environment. *J.Cell Mol. Biol.* **2**: 13-18.
- Kessler D, Diezel C, Clark DG, Colquhoun TA and Baldwin IT. 2012. Petunia flowers solve the defence/apparency dilemma of pollinator attraction by deploying complex floral blends. *Eco. Letters* **16**: 299-306
- Landry, L.G., Chapple, CCS and Last RL. 1995. *Arabidopsis* mutants lacking phenolic sunscreens exhibit enhanced ultraviolet-B injury and oxidative damage. *Pl. Physiol.***109**:1159-66
- Lattanzio V., Lattanzio V.M.T. and Cardinali A., 2006. Role of polyphenols in the resistance mechanisms of plants against fungal pathogens and insects. In : *Phytochemistry : Advances in Research* Imperato F., ed.. Trivandrum, Kerala, India
- Lindroth, R. L., Scriber, J. M., and Hsia, M. T. S. 1988. Chemical ecology of the tiger swallowtail: Mediation of host use by phenolic glycosides. *Ecology*, **69** : 814-22.
- Lovely, D.R. 2003. Cleaning up with genomics: applying molecular biology of self-bioremediation. *Nat. Rev. Microbiol.* **1**:35-44.
- M. R. Kant M.R., W. Jonckheere W., Knecht B., Lemos, F., Liu, J and Schimmel, B.C. 2015. Mechanisms and ecological consequences of plant defence induction and suppression in herbivore communities. *Ann. Bot.* **115** : 1015-51
- Mattila, P., Pihlava, J., Pihlava, M. and Hellstrom, J., 2005. 'Contents of phenolic acid, alkyl- and alkenylresorcinols and avenanthramides in commercial grain products'. *J Agric. Food Chem.* **53**: 8290.
- Naczki, M. and Shahidi, F., 2004. Extraction and analysis of phenolics in food. *J.Chromatography*, **1054**: 95'
- Ozyigit, I.I. 2007. Phenolic changes during in vitro organogenesis of cotton (*Gossypium hirsutum* L.) shoot tips. *Afr. J. Biotech.* . **7** (8):1145-1150.
- Palmer, M.A., Bernhardt, E. and Chornesky, E., 2004. Ecology for a crowded planet. *Science*, **304**: 1251-52
- Paschold A, Halitschke, R. and Baldwin, IT. 2007. Coordinating defenses: NaCOI1 mediates herbivore-induced resistance in *Nicotiana attenuata* and reveals the role of herbivore movement in avoiding defenses. *Pl. J.*, **51**: 79-91
- Perkins, LE, Cribb, BW, and Brewer, P.B. 2013. Generalist insects behave in a jasmonate-dependent manner on their host plants, leaving induced areas quickly and staying longer on distant parts. *Proc. Royal Soc. Biol. Sci.*, **280**: 20122646
- Pourcel L, Routaboul, J.M., Cheynier, V, Lepiniec L, and Debeaujon I, 2007. Flavonoid oxidation in plants: from biochemical properties to physiological functions. *Trends Pl. Sci.* **12**: 29-36
- Quideau, S., Deffieux, D., Douat-Casassus, C., and Pouységú, L. 2011. Plant polyphenols: chemical properties, biological activities, and synthesis. *Angew. Chem. Int. Ed.* **50**:586-621.
- Rodrigues E. G., Dobroff A. S. S., Cavarsan C. F., Paschoalin T., Nimrichter L., and Mortara R. A., 2010. Effective topical treatment of subcutaneous murine B16F10-Nex2 melanoma by the antimicrobial peptide gomesin. *Neoplasia*, **10**, 61-68.
- Samac, D.A. and Graham, M.A. 2007. Recent advances in legume–microbe interactions: recognition, defense response, and symbiosis from a genomic perspective. *Pl. Physiol.* **144**: 582-87.
- Seneviratne, G. and Jayasinghearachchi, H.S. 2003. Mycelial colonization by bradyrhizobia and azorhizobia. *J. Biosci.* **28**: 243-47
- Shroff R, Vergara F, and Muck A.2008. Nonuniform distribution of glucosinolates in *Arabidopsis thaliana* leaves has important consequences for plant defense. *Proc. Nat. Acad. Sci., USA* **105**: 6196-6201.
- Steidle, A., Sigl, K., Schuegger, R., Ihring, A., Schmid, M., Gantner, S., Stoffels, M., Riedel, K., Givskov M., Hartmann, A., Langebartels, A. and Eberl, L. 2001. Visualization of N-acylhomoserine lactone-mediated cell–cell communication between bacteria colonizing the tomato rhizosphere. *Appl. Env. Microbiol.* **67**, 5761-5770
- Van der Meer IM, Stam ME, Van Tunen AJ, Mol JN, and Stuitje AR. 1992. Antisense inhibition of flavonoid biosynthesis in *Petunia* anthers results in male sterility. *Pl. Cell* **4** 253-62
- Yao, L.,H., Jian, Y., M., Shi, Jian, J., Tomas-Barberan, F., A., Datta, N., Singanusong, R., and Chen, S., S. 2004. Flavonoids in food and their health benefits. *Pl. Foods Human Nutr.* **59**, pp 113.