



## Seed yield and quality of coriander (*Coriandrum sativum* L.) as influenced by seed priming

A. SARKAR, A. B. SHARANGI, <sup>1</sup>S. SOUJANNYA AND <sup>1</sup>A. DATTA

Department of Plantation, Spices, Medicinal and Aromatic Crops, <sup>1</sup>Department of Seed Science and Technology,  
Bidhan Chandra KrishiViswavidyalaya, Mohanpur - 741 252, Nadia, West Bengal

Received : 10.01.2019 ; Revised : 13.04.2020 ; Accepted : 20.04.2020

DOI : 10.22271/09746315.2020.v16.i1.1271

### ABSTRACT

Seed priming, a controlled hydration process, is a simple and cheap technology to the farmers for better germination of seed ensuring a good yield and quality. An experiment was conducted to evaluate the effects of different priming media, and duration on seed germination, growth, yield and essential oil content of coriander at the Horticultural Research Station, Mondouri, West Bengal. There were three different concentrations of chemical ( $KH_2PO_4$ ) viz., 3-, 2-, 1% and control (distilled water) with three different duration of soaking viz., 12-, 16- and 20 hrs. Seed priming with  $KH_2PO_4$  has significant influence on plant height of coriander. Maximum number of primary branches per plant (7.0), number of secondary branches per plant (12.3), maximum germination (91%), were recorded with 1%  $KH_2PO_4$  for 20 hrs. 3%  $KH_2PO_4$  for 12 hrs was found to produce maximum test weight (13.0 g), whereas, seed priming with 1%  $KH_2PO_4$  for 20 hrs out-yielded all other treatments resulting in the highest number of umbels per plant (23.3), umbellets per umbel (5.7), seed per umbel (38.4), seed yield per plant (3.5 g), seed yield per meter square (15.0 g), projected seed yield ( $13.5 \text{ q ha}^{-1}$ ), respectively. It may be concluded from the results that to obtain higher germination, seed yield and essential oil of coriander, seeds should be treated with 2%  $KH_2PO_4$  as seed priming chemical for 20 hrs under this Gangetic alluvial region as revealed from the result.

**Keywords:** Coriander, essential oil seed, germination, priming, yield

Coriander (*Coriandrum sativum* L.), is an important winter season seed spice of the family Apiaceae. Its leaves are extensively used for flavouring dishes and seeds as spices and condiments throughout the world including India. Coriander is native to South Europe and the Mediterranean region, and is extensively grown in Russia, Bulgaria, Mexico, USA, Argentina, China, Romania, Italy, Japan, Hungary, Poland, Czech, Morocco and India since human antiquity. The seeds and different parts of the plant have been reported for multiple health functions and biological activities (Nadkarni, 1976, Saeed and Tariq, 2007; Matasyoh *et al.*, 2009; Begnami *et al.*, 2010). It is used in the preparation of many household medicines to cure cold, seasonal fever, nausea, vomiting and stomach disorder. Pharmaceutical use of coriander is essentially to mask the taste of other medicinal compounds or to calm the irritating effects on the stomach that some medicines cause. Coriander leaves and seeds are valued as food mainly for its high Vitamin A and Vitamin C. Coriander oil is a clear, colourless to light yellow liquid. Indian coriander seeds are poor in oil content (0.1–0.4%). The content of essential oil in ripe fruit is comparatively low (typically, less than 1%) (Kumar *et al.*, 1977).

Seed priming is a controlled hydration process followed by re-drying that allows seed to imbibe water and begin internal biological processes necessary for germination, but which does not allow the seed to actually germinate. Priming allows some of the metabolic

processes necessary for germination to occur. In priming, seeds are soaked in different solutions with high osmotic potential. This prevents the seeds from absorbing enough water for radical protrusion, thus suspending the seeds in the lag phase (Taylor *et al.*, 1998). Seed priming has been commonly used to reduce the time between seed sowing and seedling emergence and to synchronize emergence (Parera and Cantliffe, 1994). Several investigations (Taylor and Harman 1990, Van Hulten *et al.*, 2006) confirmed that seed priming has many benefits including early and rapid emergence, stand establishment, crop establishment, higher water use efficiency, deeper roots, increase in root growth, uniformity in emergence, synchronize the germination of individual seeds break of seed dormancy, initiation of reproductive organs, better competition with weed, early flowering and maturity, resistance to environmental stresses (such as drought and salinity) and diseases (*Sclerotium rolfsii* L.) as well as increment in yield thereby providing a simple and cheap technology to the farmers for better crop production. Seed priming can be accomplished through different methods such as hydro-priming (soaking in distilled water), osmo-priming (soaking in osmotic solutions such as poly ethylene glycol, potassium salts, *e.g.*, KCl,  $K_2SO_4$ ) and plant growth inducers (CCC, Ethephon, IAA) (Capron *et al.*, 2000). Osmo-priming and hydro-priming of seeds may improve germination and emergence (Ashraf and Abu-Shakra, 1978) and may promote vigorous root growth (Carceller and Soriano, 1972).

The information regarding safe limits of priming duration and concentration in coriander and the associated effects on growth, yield and quality is very much scanty especially under Gangetic alluvial region. There are also no comprehensive understanding on germination behavior of the crop leading to field establishment and its subsequent translation towards yield and quality. Keeping in view the above, the present experiment was undertaken to evaluate the effects of different priming media and duration on seed germination on growth, yield and essential oil content of coriander.

## MATERIALS AND METHODS

An investigation on the effect of seed priming in coriander (*Coriandrum sativum* L.) was carried out at the Horticultural Research Station, Mondouri, Faculty of Horticulture, Bidhan Chandra Krishi Viswavidyalaya, Nadia, West Bengal during the year 2018-19 in the month of November to March for identifying the best seed priming concentration and duration to get the improved germination of coriander and its growth yield and quality under Gangetic alluvial plains of West Bengal. The Research Station was located at 23.5° North Latitude, 89° East Longitude having an average altitude of 9.75 m above mean sea level. The experimental site was located in subtropical humid climate with Gangetic alluvial soil having sandy clay loam texture, with good water holding capacity, well drained, and with acidic to neutral reaction (pH 6.7) and moderate fertility status (organic carbon: 0.74%, available N: 140.5 kg ha<sup>-1</sup>, available P<sub>2</sub>O<sub>5</sub>: 28.5 kg ha<sup>-1</sup>, available K<sub>2</sub>O : 87.0 kg ha<sup>-1</sup>).

The present investigation was laid out in Randomized Block Design replicated thrice with three different concentrations of potassium dihydrogen phosphate (KH<sub>2</sub>PO<sub>4</sub>) viz., C<sub>1</sub> (Seed priming with 3% KH<sub>2</sub>PO<sub>4</sub>), C<sub>2</sub> (2% KH<sub>2</sub>PO<sub>4</sub>), C<sub>3</sub> (1% KH<sub>2</sub>PO<sub>4</sub>) and one control (C<sub>0</sub>, without KH<sub>2</sub>PO<sub>4</sub>, with water). The coriander seeds were soaked for three different durations namely D<sub>1</sub> (12 hrs), D<sub>2</sub> (16 hrs), D<sub>3</sub> (20 hrs). There were twelve number of treatment combination randomized in the plots of 2 x 1.5 m<sup>2</sup> size with spacing of 25 cm X 10 cm and 120 plants per plot. The statistical analysis was done by using OP-STAT, SPSS (IBM v. 20.0) and MS Excel software.

## RESULTS AND DISCUSSION

### Seed germination, root and shoot length

A remarkable positive effect on germination of coriander seed was found on seed priming with KH<sub>2</sub>PO<sub>4</sub> as observed in fig.1. The highest germination per plant (91%) was recorded with C<sub>2</sub>D<sub>3</sub> treatment combination

(seed priming with 2% KH<sub>2</sub>PO<sub>4</sub> for 20 hrs) followed by C<sub>2</sub>D<sub>2</sub> (89.7%), C<sub>2</sub>D<sub>1</sub> (87%) and the lowest germination per plant was observed in C<sub>0</sub>D<sub>3</sub> (73.3%). The highest root and shoot length per plant (21.4 and 12.3 cm, respectively) was found with C<sub>2</sub>D<sub>3</sub> (seed priming with 2% KH<sub>2</sub>PO<sub>4</sub> for 20 hrs) treatment combination which was statistically at par with the other treatments (Table 1). The highest seedling vigour index (3000) was obtained from the treatment (C<sub>2</sub>D<sub>3</sub>) where seed priming was done with 2% KH<sub>2</sub>PO<sub>4</sub> for 20 hrs closely followed by C<sub>2</sub>D<sub>2</sub> and C<sub>3</sub>D<sub>1</sub> which were statistically at par (Fig.2). There are several other reports that seed priming can homogenize seed germination in a short period of time (Zhu, 2002; Khajeh-Hosseini, 2003)

### Morphological characters

Seed priming with KH<sub>2</sub>PO<sub>4</sub> has significant influence on plant height of coriander (Fig. 3). The observations recorded on 30, 60 and 120 days after sowing (DAS) showed that among the treatment C<sub>2</sub>D<sub>3</sub> (seed priming with 2% KH<sub>2</sub>PO<sub>4</sub> for 20 hrs) recorded the highest plant height (64.3 cm) on 30 DAS and at harvest. However, for 60 DAS C<sub>2</sub>D<sub>2</sub> treatment combination recorded the highest plant height (60.3 cm). Plants from primed seeds might enhance water and nutrients intake resulting in increased plant height as was observed by Farooq *et al.* (2006) in rice seed. The number of branches per plant has significant effect on the seed priming with KH<sub>2</sub>PO<sub>4</sub> in coriander (Fig. 4). The highest number of primary and secondary branches per plant (7 and 12.3, respectively) was recorded with C<sub>2</sub>D<sub>3</sub> (seed priming with 2% KH<sub>2</sub>PO<sub>4</sub> for 20 hrs) treatment combination. The similar result was also observed by Ranjita Devi and Sharangi (2019) in case of a few germplasms on plant height (64.8 cm) and number of primary branches (3.8 to 10.5) in coriander.

### Yield and yield attributes

The maximum number of umbels per plant<sup>-1</sup> (23.3), number of umbellets umbel<sup>-1</sup> (5.7), and seed umbel<sup>-1</sup> (38.4) (Fig 5) were recorded with C<sub>2</sub>D<sub>3</sub> (seed priming with 2% KH<sub>2</sub>PO<sub>4</sub> for 20 hrs) treatment combination. The maximum test weight (1000 seed, 13 g) was obtained from C<sub>1</sub>D<sub>1</sub> (seed priming with 3% KH<sub>2</sub>PO<sub>4</sub> for 12 hrs) followed by C<sub>3</sub>D<sub>2</sub> (12.5 g) and C<sub>2</sub>D<sub>1</sub> (12.1 g). The maximum seed yield (15.0 g m<sup>2</sup>) was obtained with C<sub>2</sub>D<sub>3</sub> (seed priming with 2% KH<sub>2</sub>PO<sub>4</sub> for 20 hrs) followed by C<sub>3</sub>D<sub>1</sub> and C<sub>2</sub>D<sub>2</sub> (14.7 g/m<sup>2</sup>) which were statistically at par and the lowest seed yield was observed in C<sub>0</sub>D<sub>3</sub> (12.4g/m<sup>2</sup>). The seed yield (g plant<sup>-1</sup>) has also a similar trend with significant effects on the seed priming with KH<sub>2</sub>PO<sub>4</sub> in coriander. The seed priming with KH<sub>2</sub>PO<sub>4</sub> has a significant effect on projected seed yield. The highest projected seed yield (13.5 q/ha) was recorded with C<sub>2</sub>D<sub>3</sub>

**Table1: Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on root and shoot length of coriander**

Treatments	Root length (cm)*	Shoot length (cm)*
C <sub>1</sub> D <sub>1</sub> (Seed priming with 3% $\text{KH}_2\text{PO}_4$ for 12 hrs)	18.5±0.38	9.8±0.43
C <sub>1</sub> D <sub>2</sub> (Seed priming with 3% $\text{KH}_2\text{PO}_4$ for 16 hrs)	19.5±0.35	9.9±0.18
C <sub>1</sub> D <sub>3</sub> (Seed priming with 3% $\text{KH}_2\text{PO}_4$ for 20 hrs)	20.4±0.61	10.2±0.31
C <sub>2</sub> D <sub>1</sub> (Seed priming with 2% $\text{KH}_2\text{PO}_4$ for 12 hrs)	20.1±0.17	11.1±0.35
C <sub>2</sub> D <sub>2</sub> (Seed priming with 2% $\text{KH}_2\text{PO}_4$ for 16 hrs)	19.9±0.22	10.9±0.32
C <sub>2</sub> D <sub>3</sub> (Seed priming with 2% $\text{KH}_2\text{PO}_4$ for 20 hrs)	21.4±0.38	12.3±0.21
C <sub>3</sub> D <sub>1</sub> (Seed priming with 1% $\text{KH}_2\text{PO}_4$ for 12 hrs)	20.7±0.78	11.2±0.49
C <sub>3</sub> D <sub>2</sub> (Seed priming with 1% $\text{KH}_2\text{PO}_4$ for 16 hrs)	19.1±0.49	10.8±0.23
C <sub>3</sub> D <sub>3</sub> (Seed priming with 1% $\text{KH}_2\text{PO}_4$ for 20 hrs)	19.4±0.38	10.7±0.58
C <sub>0</sub> D <sub>1</sub> (Distilled water for 12 hrs)	18.2±0.58	11.2±0.23
C <sub>0</sub> D <sub>2</sub> (Distilled water for 16 hrs)	17.7±0.35	10.5±0.93
C <sub>0</sub> D <sub>3</sub> (Distilled water for 20 hrs)	17.6±0.33	10.2±0.20
<b>SEm(±)</b>	<b>0.46</b>	<b>0.42</b>
<b>LSD(0.05)</b>	<b>1.36</b>	<b>1.24</b>

\*Values represent Mean±SE

**Table 2 : Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on seed yield of coriander**


Treatments	Seed Yield (g plant <sup>-1</sup> )	Seed Yield (g m <sup>2</sup> )	Projected Seed Yield(q ha <sup>-1</sup> )
C <sub>1</sub> D <sub>1</sub>	3.1	13.5	12.2
C <sub>1</sub> D <sub>2</sub>	3.2	13.9	12.5
C <sub>1</sub> D <sub>3</sub>	3.2	14.1	12.7
C <sub>2</sub> D <sub>1</sub>	3.3	14.4	12.9
C <sub>2</sub> D <sub>2</sub>	3.4	14.7	13.2
C <sub>2</sub> D <sub>3</sub>	3.5	15.0	13.5
C <sub>3</sub> D <sub>1</sub>	3.4	14.7	13.2
C <sub>3</sub> D <sub>2</sub>	3.1	13.5	12.2
C <sub>3</sub> D <sub>3</sub>	3.3	14.1	12.7
C <sub>0</sub> D <sub>1</sub>	3.0	13.1	11.8
C <sub>0</sub> D <sub>2</sub>	2.9	12.7	11.5
C <sub>0</sub> D <sub>3</sub>	2.9	12.4	11.2
<b>SEm(±)</b>	<b>0.16</b>	<b>0.51</b>	<b>0.12</b>
<b>LSD(0.05)</b>	<b>0.47</b>	<b>1.52</b>	<b>0.36</b>

**Table 3 : Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on essential oil (%) of coriander**


Treatments	Essential oil content (%) of seed*
C <sub>1</sub> D <sub>1</sub> (Seed priming with 3% $\text{KH}_2\text{PO}_4$ for 12 hrs)	0.26±0.003
C <sub>1</sub> D <sub>2</sub> (Seed priming with 3% $\text{KH}_2\text{PO}_4$ for 16 hrs)	0.25±0.003
C <sub>1</sub> D <sub>3</sub> (Seed priming with 3% $\text{KH}_2\text{PO}_4$ for 20 hrs)	0.26±0.003
C <sub>2</sub> D <sub>1</sub> (Seed priming with 2% $\text{KH}_2\text{PO}_4$ for 12 hrs)	0.27±0.003
C <sub>2</sub> D <sub>2</sub> (Seed priming with 2% $\text{KH}_2\text{PO}_4$ for 16 hrs)	0.27±0.007
C <sub>2</sub> D <sub>3</sub> (Seed priming with 2% $\text{KH}_2\text{PO}_4$ for 20 hrs)	0.28±0.006
C <sub>3</sub> D <sub>1</sub> (Seed priming with 1% $\text{KH}_2\text{PO}_4$ for 12 hrs)	0.26±0.003
C <sub>3</sub> D <sub>2</sub> (Seed priming with 1% $\text{KH}_2\text{PO}_4$ for 16 hrs)	0.26±0.003
C <sub>3</sub> D <sub>3</sub> (Seed priming with 1% $\text{KH}_2\text{PO}_4$ for 20 hrs)	0.25±0.003
C <sub>0</sub> D <sub>1</sub> (Distilled water for 12 hrs)	0.23±0.003
C <sub>0</sub> D <sub>2</sub> (Distilled water for 16 hrs)	0.24±0.003
C <sub>0</sub> D <sub>3</sub> (Distilled water for 20 hrs)	0.23±0.003
<b>SEm(±)</b>	<b>0.004</b>
<b>LSD(0.05)</b>	<b>0.011</b>

\*Values represent Mean±SE


## Seed yield and quality of coriander




**Fig. 1:** Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on germination of coriander




**Fig. 2:** Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on seedling vigour index of coriander




**Fig. 3:** Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on plant height of coriander



**Fig. 4:** Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on primary and secondary branches of coriander



**Fig. 5:** Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on yield attributing parameters of coriander



**Fig. 6:** Effect of seed priming with  $\text{KH}_2\text{PO}_4$  on essential oil of coriander

(seed priming with 2%  $\text{KH}_2\text{PO}_4$  for 20 hrs) treatment combination followed by  $\text{C}_3\text{D}_1$  ( $13.2 \text{ q ha}^{-1}$ ),  $\text{C}_2\text{D}_2$  ( $13.2 \text{ q ha}^{-1}$ ) and the lowest seed yield quintal per hectare was observed in  $\text{C}_0\text{D}_3$  ( $11.2 \text{ q ha}^{-1}$ ). Seed priming might have increased the performance by altering the mechanism of enzymes which are responsible for enhancing yield attributes and yield thereby (Jamshidian and Talat, 2017).

### Essential oil content (%)

It is evident from table 3 and figure 6 that seed priming with  $\text{KH}_2\text{PO}_4$  has significant effect on essential oil content in coriander seed. The essential oil content (0.28 %) was the highest as recorded with  $\text{C}_2\text{D}_3$  (seed priming with 2%  $\text{KH}_2\text{PO}_4$  for 20 hrs) followed by  $\text{C}_2\text{D}_2$  (0.27 %) and  $\text{C}_2\text{D}_1$  (0.27 %) whereas the lowest oil content was observed in  $\text{C}_0\text{D}_3$  (0.23 %). The essential oil content and composition of *C. sativum* can be influenced by cultivation practices, ontogenetic and genetic factors (Msaada *et al.*, 2007; Telci *et al.*, 2006). The essential

oil content of coriander in the present study corresponded with the earlier reports obtained by Bandara *et al.* (2000).

The study showed a significant effect of both concentration of seed priming chemical and duration of soaking on germination, growth, yield and quality of coriander. So it may be concluded from the results that to obtain higher germination, seed yield and essential oil of coriander, seeds should be treated with 2%  $\text{KH}_2\text{PO}_4$  as seed priming chemical for 20 hrs under this Gangetic alluvial region.

### REFERENCES

- Ashraf, C. M. and Abu-Shakra, S. 1978. Wheat seed germination under low temperature and moisture stress. *Agronomy J.*, **70**: 135-39.
- Bandara, M., Wildschut, C., Russel, E., Ost, L., Simo, T., and Weber, J. 2000. Special crops program (Brooks). Alberta, Agriculture, Food, and Rural Development. Crop Diversification Centres

- Annual Report. Alberta, Canada. Available at: [http://www.agric.gov.ab.ca/ministry/pid/cdc/00/sc\\_brooks.html](http://www.agric.gov.ab.ca/ministry/pid/cdc/00/sc_brooks.html).
- Begnam AF, Duarte MCT, Furletti RV. 2010. Antimicrobial potential of *Coriandrum sativum* L. against different candida species in vitro. *Food Chem.*, **118**: 74-77.
- Carceller, M. S. and Soriano, A. 1972. Effect of treatments given to grain, on the growth of wheat roots under drought conditions. *Can. J. Bot.*, **50**: 105-08.
- Capron, I., Corbineau, F., Dacher, F., Job, C., Côme, D. and Job, D. 2000. Sugarbeet seed priming: effects of priming conditions on germination, solubilization of 11-S globulin and accumulation of LEA proteins. *Seed Sci. Res.*, **10**(3): 243-54.
- Farooq, M., Basra, S.M.A., Warraich, E.A. 2006. Optimization of hydro-priming techniques for rice seed invigoration. *Seed Sci. Technol.*, **34** (2): 507-12.
- Jamshidian, Z. and Talat, F. 2017. Effects of seed priming on morphological and phonological characteristics of the coriander (*Coriandrum sativum* L.). *Adv. Pl. Agric. Res.*, **7**(6): 411-15.
- Khajeh-Hosseini, M., Powell, A.A., Bringham, I.J. 2003. The interaction between salinity stress and seed vigour during germination of soybean seeds. *Seed Sci. Technol.*, **31**: 715-25.
- Kumar, C. R., Sarwar, M. and Dimri, B. P. 1977. Bulgarian coriander in India and its future prospects in export trade. *Indian perfumer*, **21**: 146-50.
- Matasyoh, JC, Maiyo, ZC, Ngure, R. and Chepkorir, R. 2009. Chemical composition and antimicrobial activity of essential oil of *Coriandrum sativum*. *Food Chem.*, **113**: 526-29.
- Msaada, K.H.K., Ben Taarit, M., Chahed, T., Kchouk, M.E. and Marzouk, B. 2007. Changes on essential oil composition of coriander (*Coriandrum sativum* L.) fruits during three stages of maturity. *Food Chemistry*, **102** : 1131- 34.
- Nadkarni, A.K., 1976. Indian Materia Medica Popular. Prakshan, Bombay, p. 381.
- Parera, C. A. and Cantliffe, D. J. 1994. Pre-sowing seed priming. *Hortic. Rev.*, **16**: 109-41.
- Ranjita Devi, A. and Sharangi, A.B. 2019. Morphological character and seed yield potential of coriander genotypes under Gangetic Alluvial region of West Bengal. *Int. J. Curr. Microbiol. App. Sci.*, **8**(4): 775-82.
- Saeed S, Tariq P. 2007. Antimicrobial activities of *Emblica officinalis* and *Coriandrum sativum* against Gram-positive bacteria and *Candida albicans*. *Pak. J. Bot.*, **39**: 913-17.
- Said-Al Ahl, H.A.H., El Gendy, A.G. and Omer, E.A. 2014. Effect of Ascorbic Acid, Salicylic Acid on Coriander Productivity and Essential Oil Cultivated in two Different Locations. *Advances in Env. Biol.*, **8**(7): 2236-50.
- Taylor, A.G., Harman, G.E. (1990) Concepts and technologies of selected seed treatments. *Ann. Rev. Phytopathol.*, **28**: 321-29.
- Taylor, A. G., Allen, P. S., Bennett, M. A., Bradford, K. J., Burris, J. S. and Misra, M. K. 1998. Seed enhancements. *Seed Sci. Res.*, **8**(2): 245-5.
- Telci, I., Toncer, O.G., and Sahbaz, N. 2006. Yield, essential oil content and composition of *Coriandrum sativum* varieties (var. *vulgare* Alef and var. *microcarpum* DC.) grown in two different locations. *J. Essential Oil Res.*, **18**: 189-93.
- Van Hulst, M., Pelser, M., Van Loon, L.C., Pieterse, C.M.J. and Ton, J. 2006. Costs and benefits of priming for defense in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA*, **103**: 5602-07.
- Zhu, J.K. 2002. Salt and drought stress signal transduction in plants. *Annu. Rev. Plant Biol.*, **53**: 247-73.