

## Growth and yield parameters of okra (*Abelmoschus esculentus*) influenced by Diazotrophs and chemical fertilizers

B. MAL, P. MAHAPATRA, S. MOHANTY AND H. N. MISHRA

Department of Vegetable Science, Department of Soil Science and Agricultural Chemistry  
Orissa University of Agriculture and Technology  
Bhubaneswar, Odisha, India

Received: 9-10-2013, Revised: 18-11-2013, Accepted: 20-11-2013

### ABSTRACT

Field experiments were conducted during the summer seasons of 2010 and 2011 in College of Agriculture, Orissa University of Agriculture and Technology (OUAT), Bhubaneswar, to assess the effect of diazotrophs (Biofertilizers - *Azotobacter*, *Azospirillum*, Phosphate Solubilizing Bacteria) and chemical fertilizers along with vermicompost on okra cv Mahyco-10 in terms of growth and yield. The experiment consisted of nine treatments including the control and replicated thrice in randomized block design. The maximum plant height (148.97 cm), leaf area (434.99 cm<sup>2</sup>), number of nodes (30.16), fruit length (16.45 cm), fruit girth (1.62 cm), single fruit weight (18.70 g) and plant biomass-fresh weight (548.74 q ha<sup>-1</sup>), were observed in the treatment receiving FYM@10 t ha<sup>-1</sup> + 100% NPK + vermicompost @ 5 t ha<sup>-1</sup> along with biofertilizers. The maximum number of fruits per plant was recorded with T<sub>9</sub> (FYM@10 t ha<sup>-1</sup> + 75% NPK + vermicompost@5 t ha<sup>-1</sup> + biofertilizers). Application of higher dose of fertilizers increased the fruit yield of okra considerably, where the yield varied between 80.24 q ha<sup>-1</sup> to 228.38 q ha<sup>-1</sup>. The study led to a conclusion, that the maximum growth parameters, highest yield and yield attributing characters of okra could be achieved by integrated use of diazotrophs, vermicompost and chemical fertilizers.

**Key words:** Diazotrophs, FYM, okra, vermicompost and yield

Okra is an important vegetable crop which supplies higher nutrition (carbohydrates, fats, protein, minerals and vitamins) in our diet. The main challenge before India is to increase the production of quality food in a sustainable manner and feeding the country's large population and increasing the income of the farmer. The requirements of fertilizers in okra are important for the early growth and total production of fruit yield. Integrated use of organic and inorganic fertilizers can improve crop productivity (Satyanarayana *et al.*, 2002). The soil enriched with vermicompost provides additional substances that are not found in chemical fertilizers (Kale, 1998). Naidu *et al.* (2000) concluded that significant increase of microbes in soil was found with application of manures, vermicompost and biofertilizers. Inoculation with diazotrophs (*Azospirillum*, *Azotobacter* and PSB) in okra helped fixing atmospheric N, increased phosphate availability produced growth promoting and antifungal substances and finally increased the total yield. The objective of the study was to assess the effect of FYM, chemical fertilizers, vermicompost along with biofertilizers on okra in terms of growth and yield.

### MATERIALS AND METHODS

Field experiment was conducted to evaluate the performances of okra variety *i.e.*, F1 hybrid (Mahyco-10) under different nutrient management practices. The mechanical composition were recorded as sand (84%), silt clay (16%) and clay (11%) and chemical composition as soil pH (5.34), available soil nitrogen (180 kg ha<sup>-1</sup>), available soil phosphorus (20 kg ha<sup>-1</sup>) and available soil potassium (120 kg ha<sup>-1</sup>).

The total precipitations during the cropping period recorded from February to June 2010 and 2011 were 190.9 mm and 249.1 mm respectively. The maximum and minimum temperatures during the first and second cropping year (2010) varied from 33.4°C to 39.2°C and 18.4°C to 24.3°C respectively.

The experiment was laid out in randomized block design with 3 replications. Altogether 27 plots of 9 m<sup>2</sup> each were prepared for the experiment. FYM@10 t ha<sup>-1</sup> was applied in all the treatments as basal dose. The recommended N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O fertilizer doses for F<sub>1</sub> Hybrid okra *i.e.* MAHYCO-10 was 200,100,100 kg ha<sup>-1</sup>. Fertilizers were applied in split doses following during the cropping period. Vermicompost (5 t ha<sup>-1</sup> and 2.5 t ha<sup>-1</sup>) was also applied to the plot as per the requirement of the treatment. Further mixed culture of bio-fertilizers *i.e.* *Azotobacter*, *Azospirillum* and PSB (1:1:1) was used before sowing as per the treatments. The experiment consisted of nine treatments including the control treatment *viz.*, T<sub>1</sub> - Control (FYM @10 t ha<sup>-1</sup>), T<sub>2</sub> - FYM (10 t ha<sup>-1</sup>) + 100 % RDF (NPK), T<sub>3</sub> - FYM (10 t ha<sup>-1</sup>) + 75 % RDF(NPK), T<sub>4</sub> - FYM (10 t ha<sup>-1</sup>) + 100 % RDF (NPK) + Bio-fertilizers [*Azotobacter*, *Azospirillum* and PSB (1:1:1) @6 kg ha<sup>-1</sup>], T<sub>5</sub> - FYM (10 t ha<sup>-1</sup>) + 75 % RDF (NPK) + Bio-fertilizers [*Azotobacter*, *Azospirillum* and PSB (1:1:1) @6 kg ha<sup>-1</sup>], T<sub>6</sub> - FYM (10 t ha<sup>-1</sup>) +100 % RDF (NPK) + Vermicompost (5 t ha<sup>-1</sup>) + Bio-fertilizers [*Azotobacter*, *Azospirillum* and PSB (1:1:1) @6 kg ha<sup>-1</sup>], T<sub>7</sub> - FYM (10 t ha<sup>-1</sup>) +75 % RDF (NPK) +Vermicompost(2.5 t ha<sup>-1</sup>) + Bio-fertilizers [*Azotobacter*, *Azospirillum* and PSB (1:1:1) @6 kg

ha<sup>-1</sup>], T<sub>8</sub> - FYM (10 t ha<sup>-1</sup>) + 100% RDF (NPK) + Vermicompost (2.5 t ha<sup>-1</sup>) + Bio-fertilizers [*Azotobacter*, *Azospirillum* and PSB (1:1:1) @6 kg ha<sup>-1</sup>] and T<sub>9</sub> - FYM(10 t ha<sup>-1</sup>)+75 % RD (NPK) +Vermicompost (5 t ha<sup>-1</sup>) + Bio-fertilizers [*Azotobacter*, *Azospirillum* and PSB (1:1:1) @6 kg ha<sup>-1</sup>].

Ten plants were selected from each plot to record the biometric observations at various stages of crop growth. Growth parameters include plant height (cm), leaf area per leaf (cm<sup>2</sup>), days to emergence of first flower and number of nodes per plant etc was measured prior to harvest. Post harvest observations viz., fruit length, fruit girth, number of fruits, single fruit weight, plant biomass (fresh weight) and yield of the crop were recorded after harvesting. The mean data and physico-chemical estimates were subjected to proper statistical analysis as per Randomized Block Design system. The F-test was used for testing the significance of findings. Standard error for each factor was worked out to compare the means of two treatments. The Least Significant difference (LSD) was calculated at 5% level of significance.

## RESULTS AND DISCUSSION

### Plant height

The maximum plant height (148.97 cm) was obtained in T<sub>6</sub> when the highest amount of fertilizer (FYM@10 t ha<sup>-1</sup> +100% RDF + vermicompost @5 t ha<sup>-1</sup> + biofertilizer) dose was applied followed by FYM@10 t ha<sup>-1</sup> + 100% RDF + vermicompost @2.5 t ha<sup>-1</sup> + biofertilizer and both were statistically significant over rest of the treatments. The lowest plant height (110.90 cm) was recorded when the crop was grown with FYM only and sole application of inorganic fertilizers (100 % RDF or 75 % RDF) with FYM which resulted in poor performance as compared to the integrated application of fertilizers. It might be due to the increase in the nutrient availability and preponderance of different groups of microorganisms in soil, which create a favourable condition for proper vegetative growth in general and increased plant height in particular. The highest dose of nitrogen might have enhanced cell division and formation of more tissues resulting in luxuriant vegetative growth and thereby increasing plant height (Meyer and Anderson, 2003).

### Leaf area

Significant differences were exhibited among different treatments with respect to leaf area. The highest (434.99 cm<sup>2</sup>) and lowest (303.81 cm<sup>2</sup>) leaf area were found in T<sub>6</sub> and T<sub>1</sub> respectively. From the present investigation, it is seen that there was a significant effect of application of FYM, 100% NPK along with vermicompost and biofertilizers on increasing the leaf area as compared to the control.

The large leaf area might have contributed to the higher leaf dry weight grown with higher dose of potassium.

### Number of nodes

The highest number of nodes (30.16) per plant was recorded in T<sub>6</sub> and the lowest (16.89) was in T<sub>1</sub>. In okra, each node is a very important character for producing fruit. Application of FYM@10 t ha<sup>-1</sup> + 100% RDF + vermicompost @5 t ha<sup>-1</sup> and biofertilizer significantly increased the number of nodes per plant. This might be due to the better availability and uptake of plant nutrients, more specifically N, P and K, resulting in better photosynthesis and protein synthesis. Kumar *et al.* (2009) observed that application of 25% recommended dose of N through vermicompost significantly improved nodes per plant.

### Number of days for emergence of first flower

The minimum number of days (37.40 days) taken for emergence of the first flower was in T<sub>6</sub> and the maximum number of days (45.02 days) taken was in control. This could be due to nitrogen and other inputs like vermicompost and biofertilizers which encouraged the differentiation of bud resulting in earlier flowering. Maximum number of days was required to flower where no minimum dose of fertilizer was applied; probably due to the nutrient stress resulting in late flowering. Earliness in days to flowering in okra was observed with the integrated nutrient application (chemical fertilizers, organic manures and biofertilizers) by Prabhu *et al.* (2002).

### Number of fruits

The highest (18.36) and lowest (10.39) numbers of fruits per plant were recorded in T<sub>9</sub> and T<sub>1</sub> respectively. The numbers of fruits per plant, the most important factor of fruit yield in okra were also significantly influenced by the combined application of chemical fertilizers, vermicompost and biofertilizer as compared to control. This might be due to the better availability and uptake of nutrients by plants for a longer duration of crop growth. Similar findings of significantly higher number of fruits per plant by integrated application of fertilizers have also been reported by Prabhu *et al.* (2003) in okra.

### Fruit length and girth

The maximum (16.45 cm and 1.62 cm) and the minimum (9.99 cm and 1.39 cm) fruit length and girth were observed with highest dose of fertilizer i.e.FYM@10 t ha<sup>-1</sup> 100% RDF + vermicompost @5 t ha<sup>-1</sup> + biofertilizer. This might be attributed to the increased availability of NPK and water at the critical stages of the crop growth resulting early




Fig. 1 Maximum and minimum temperatures during cropping season

Table 1: Effect of diazotrophs and chemical fertilizers on growth parameters of okra (Pooled)

Treatments	Plant height (cm)	Leaf area (cm <sup>2</sup> )	No. of nodes plant <sup>-1</sup>	Days to emergence of 1 <sup>st</sup> flower	No. of fruits per plant <sup>-1</sup>	Fruit length (cm)	Fruit girth (cm)	Single fruit weight (g)	Fruit yield (q ha <sup>-1</sup> )	Plant biomass-fresh weight (q ha <sup>-1</sup> )
T <sub>1</sub> Control (FYM 10 t ha <sup>-1</sup> )	110.90	303.81	16.89	45.02	9.99	1.39	11.62	80.24	387.38	10.39
T <sub>2</sub> T <sub>1</sub> + 100% RDF	135.22	369.48	26.05	39.93	14.69	1.57	15.33	172.96	448.03	16.94
T <sub>3</sub> T <sub>1</sub> +75% RDF	133.66	361.82	25.27	40.07	14.46	1.56	14.75	168.64	442.93	17.17
T <sub>4</sub> T <sub>1</sub> +100% RDF + Bio-fertilizer	138.88	389.89	26.93	39.59	15.05	1.58	16.07	183.58	469.28	17.14
T <sub>5</sub> T <sub>1</sub> +75% RDF + Bio-fertilizer	137.68	380.38	26.14	39.65	14.94	1.58	16.03	179.70	461.29	16.85
T <sub>6</sub> T <sub>1</sub> + 100% RDF + Vermicompost(5 t ha <sup>-1</sup> ) + Biofertilizer	148.97	434.99	30.16	37.40	16.45	1.62	18.70	228.38	548.74	18.33
T <sub>7</sub> T <sub>1</sub> + 75% RDF +Vermicompost(2.5t ha <sup>-1</sup> ) +Biofertilizer	141.18	400.58	27.75	39.27	15.37	1.60	17.19	203.67	496.23	17.79
T <sub>8</sub> T <sub>1</sub> +100% RDF +Vermicompost(2.5t ha <sup>-1</sup> ) +Biofertilizer	144.95	416.96	29.11	38.86	15.98	1.61	17.88	211.25	507.32	17.75
T <sub>9</sub> T <sub>1</sub> +75% RDF +Vermicompost(5 t ha <sup>-1</sup> ) + Biofertilizer	142.27	408.22	28.42	38.32	16.23	1.60	17.47	213.71	517.12	18.36
<b>SEM(±)</b>	<b>0.19</b>	<b>0.85</b>	<b>0.16</b>	<b>0.16</b>	<b>0.16</b>	<b>0.01</b>	<b>0.12</b>	<b>0.713</b>	<b>0.29</b>	<b>0.14</b>
<b>LSD(0.05)</b>	<b>0.56</b>	<b>2.44</b>	<b>0.45</b>	<b>0.45</b>	<b>0.46</b>	<b>0.02</b>	<b>0.34</b>	<b>2.04</b>	<b>0.83</b>	<b>0.39</b>

Note: RDF - Recommended dose of fertilizer

establishment, vigorous growth and development of plants leading to longer and wider fruits. Higher value in fruit girth of okra observed due to integrated application of application of fertilizers by Naidu *et al.* (2002).

#### Fruit weight

The maximum (18.70 g) and the minimum (11.62 g) weight of the single fruit increases with treatment T<sub>6</sub> and T<sub>1</sub> respectively. This might occur due to increased photosynthetic area and translocation

of photosynthates in plants which subsequently accelerated the formation of more number of large sized fruits with more number of seeds per fruits resulting in increase in fruit weight.

### Fruit yield

The maximum yield (228.38 q ha<sup>-1</sup>) was recorded in T<sub>6</sub> which was significantly superior to rest of the treatments tried in the experiment. However, as expected the lowest yield (80.24 q ha<sup>-1</sup>) was recorded in T<sub>1</sub>. Jayapandi and Balakrishnan (1990) made a correlation study of yield components in okra and reported that characters like days to flowering, plant height, nodes per plant, fruit length, fruit weight and fruits per plants were significantly and positively correlated with the yield. Use of biofertilizers can replace the application of 75% of the recommended dose of nitrogen and phosphorus chemical fertilizers (El-Shaikh, 2005). It is evident from the experiment that the enhancement in plant growth attributes by the application of FYM@10 t ha<sup>-1</sup>, nitrogenous fertilizer as 100% of the needed requirements plus vermicompost @5 t ha<sup>-1</sup> and the inoculation with bio-fertilizers reflected on the total pod yield. Similar results were reported earlier by Jha and Mathur (1993).

### Plant biomass

The maximum plant biomass (548.74q ha<sup>-1</sup>) were recorded in V<sub>1</sub>T<sub>6</sub> (FYM@10 t ha<sup>-1</sup> + 100% RDF + vermicompost @5 t ha<sup>-1</sup> + biofertilizer) followed by V<sub>1</sub>T<sub>9</sub> (517.12q ha<sup>-1</sup>), V<sub>1</sub>T<sub>8</sub> (507.32q ha<sup>-1</sup>) and V<sub>1</sub>T<sub>7</sub> (496.23 q ha<sup>-1</sup>). This is might be due to integrated application of nutrients with higher dose of nitrogenous fertilizers which increased the dry matter content. Shanthi and Vijayakumari (2002) recorded higher fresh plant weight due to the integrated use of NPK and neem cake at 30 days after sowing in okra cv Arka Anamika.

The present investigation reveals that the maximum vegetative growth, yield attributing characters and fruit yield was found in integrated application of fertilizers as compared to recommended dosage of chemical fertilizers and organic manure alone. It is due to the effective utilization of N, P, K by the application of FYM@10 t ha<sup>-1</sup> + 100% of the recommended dose of NPK along with diazotrophs in presence of vermicompost @5 t ha<sup>-1</sup>.

### REFERENCES

- Shaikh, K.A.A. 2005. Effect of phosphorus and potassium fertilization on green yield, seed production and quality of two okra cultivars in reclaimed soils. *Assiut J. Agric. Sci.*, **36**: 51-68.
- Jayapandi, A. and Balkrishnan, R.1990. Correlation analysis in bhendi [*Abelmoschus esculentus* (L) Moench]. *South Indian Hort.* **38**:83-85.
- Jha, D. and Mathur, R. S. 1993. Combined effect of nitrogenous fertilizers and *Azospirillum brasilense* on the yield of nitrogen uptake by Pearl millet. *Int. J. Trop. Agric.*, **11**: 31-35.
- Kale R.D. 1998. *Earthworm Cinderella of Organic Farming*. Prosm Book Pvt Ltd, Bangalore, India. P.88.
- Kumar, R. P., Singh, O.N., Singh, Y., Dwivedi, S., Singh, J.P. 2009. Effect of integrated nutrient management on growth, yield, nutrient uptake and economics of french bean (*Phaseolus vulgaris*). *Indian J. Agril Sci.*, **79**: 56-61.
- Meyer, B.S. and Anderson, D.B. 2003. Nitrogen and phosphorus leaching during the decomposition of broad leaf forest litter. *Polish J. Soil Sci.* **36**, 21-29.
- Naidu, A.K.; Kushwah, S.S.; Dwivedi, Y.C.2000. Performance of organic manures, bio and chemical fertilizers and their combinations on microbial population of soil and growth and yield of Okra. *JNKVV Res. J.*, **33**: 34-38.
- Prabhu, T.; Ismail, S.; Sajindranath, A.K. and Savithiri, R. 2002a. Effect of integrated nutrient management on yield, dry matter and nutrient content in okra. *Orissa J. Hort.*, **30**: 52-56.
- Prabhu, T.; Narwadekar, P.R.; Sannindranath, A.K. and Rofi, Mohd. 2003. Effect of integrated nutrient management on growth and yield of okra (*Abelmoschus esculentus* (L.) Moench) cv. Parbhani Kranti. *Orissa J. Hort.*, **31**: 17-21.
- Satyanarayana, V., P.V.V. Prasad, V. R.K. Murthy and K. J. Boote. 2002. Influence of integrated use of farmyard manure and inorganic fertilizers on yield and yield components of irrigated lowland rice. *Indian J. Pl. Nutr.*, **25**:2081-90.
- Shanthi, N; Vijayakumari, B.2002. Effect of NPK with different organic manures on biometric parameters of bhendi (*Abelmoschus esculentus* L. var. Arka Anamika). *J. Phytological Res.*, **15**: 209-12.